

VOZOLE - PF

(Voriconazole for injection 1mg/vial – Lyophilized Powder)
Preservative Free



- Superior anti-fungal activity
- Lyophilized form enhances drug stability and solubility
- No side effects, established safety profile
- Economical pack
- Refrigeration not required

DESCRIPTION

Vozole-PF contains Voriconazole an antifungal triazole used against broad range of fungal pathogens. It is a sterile, non-pyrogenic, and available as 1 mg lyophilized powder for injection. This 1 mg lyophilized Voriconazole is reconstituted with 1ml of sterile water for injection to get the 0.1% w/v solution.

INDICATIONS

Vozole-PF should be administered primarily to patients with progressive, potentially sight threatening fungal infections. It is specifically used in the treatment of fungal keratitis caused by *Aspergillus spp.*, *Fusarium spp.* and *Candida spp.* and visually devastating disease due to *Scedosporium apiospermum*.

Clinical case reports have shown Voriconazole exhibits broad spectrum antifungal action and ocular efficacy in the cases of post cataract endophthalmitis due to *Scopulariopsis spp.*, *Aspergillus niger*, *Aspergillus flavus*, *Verticillium spp.*, and *Candida albicans* endophthalmitis with severe vitreous inflammation and subretinal infiltrates. It is also effective in the treatment of endogenous endophthalmitis due to *Scedosporium apiospermum*, *Aspergillus terreus*, *Scedosporium apiospermum*, *Fusarium* and Contact-lens related *Fusarium solani keratitis*.

WARNING AND PRECAUTIONS

Vozole-PF is recommended for intra-ocular injection and for single use only. Caution should be used in prescribing voriconazole to patients with hypersensitivity to other azoles. Concomitant use of certain drugs like Cyclosporine, Tacrolimus, Macrolide antibiotics, Terfenadine, Astemizole, Prednisolone, Benzodiazepines, Warfarin can alter the Voriconazole activity.

CONTRAINDICATIONS

Vozole-PF is contraindicated in patients with known hypersensitivity to Voriconazole or any of its excipients.

ADVERSE REACTIONS

Adverse effects including visual disturbances including abnormal vision, colour vision, change and / or photophobia are observed which typically resolve within 1 month. Elevations in hepatic enzymes levels can occur.

DOSAGE AND ADMINISTRATION

Vozole-PF sterile powder for injection is to be reconstituted with 1 ml of sterile water for injection to get 0.1% w/v. After reconstitution use the drug as directed by a registered ophthalmologist. The reconstituted solution is for single use only.

SUPPLY

Supplied as 1mg/vial in a sterile lyophilized powder form.

STORAGE

Store at a temperature between 2°C and 35°C.

Information published in this catalogue is subject to change without notification.

No. 1, Sivagangai Main Road, Veerapanjan, Madurai 625 020, India

Phone: 91 452 2446100 Fax: 91 452 2446200

E-mail: info@aurolab.com Web: www.aurolab.com